

The Poincaré series of the algebra of rational functions which are regular outside hyperplanes

HIROKI HORIUCHI and HIROAKI TERAOKA *

*Tokyo Metropolitan University, Mathematics Department
Minami-Ohsawa, Hachioji, Tokyo 192-0397, Japan*

Abstract

Let Δ be a finite set of nonzero linear forms in several variables with coefficients in a field $\mathbf{K}$ of characteristic zero. Consider the $\mathbf{K}$ -algebra $R(\Delta)$ of rational functions on V which are regular outside $\bigcup_{\alpha \in \Delta} \ker \alpha$. Then the ring $R(\Delta)$ is naturally doubly filtered by the degrees of denominators and of numerators. In this paper we give an explicit combinatorial formula for the Poincaré series in two variables of the associated bigraded vector space $\overline{R}(\Delta)$. This generalizes the main theorem of [8].

Mathematics Subject Classification (2000): 32S22, 13D40, 52C35

1 Introduction and main results

Let V be a vector space of dimension ℓ over a field $\mathbf{K}$ of characteristic zero. Let V^* be the dual space of V . Suppose that Δ is a finite subset of nonzero elements of V^* . We assume that no two vectors of Δ are proportional. Let $S = S(V^*)$ be the symmetric algebra of V^* . Then S may be regarded as the ring of polynomial functions on V .

Definition 1.1. *Let $R(\Delta)$ be the $\mathbf{K}$ -algebra of rational functions on V which are regular outside the set $\bigcup_{\alpha \in \Delta} \{\alpha = 0\}$. In other words, $R(\Delta) = S[\Delta^{-1}]$, where $\Delta^{-1} := \{1/\alpha \mid \alpha \in \Delta\}$.*

Let f/g be a homogeneous element of $R(\Delta)$, i.e., both f and g are homogeneous polynomials. Then the (total) **degree** of f/g is defined by $\deg(f/g) = \deg f - \deg g$. Let $R = R(\Delta)$, $p, q \in \mathbb{Z}$. Define a $\mathbf{K}$ -vector subspace

$$R_q^p := \left\langle \frac{f}{g} \in R \mid \deg f \leq p, \deg g \leq q \right\rangle_{\mathbf{K}}$$

*partially supported by the Grant-in-aid for scientific research (No.1144002), the Ministry of Education, Sports, Science and Technology, Japan

of R . Here the notation $\langle T \rangle_{\mathbf{K}}$ stands for the $\mathbf{K}$ -vector subspace of R spanned by a subset T of R . Agree that $R_q^p = 0$ if either $p < 0$ or $q < 0$. Then we have a double filtration on R :

$$\cdots \supseteq R_q^p \supseteq R_q^{p-1} \supseteq \cdots \quad \text{and} \quad \cdots \supseteq R_q^p \supseteq R_{q-1}^p \supseteq \cdots$$

Define

$$\overline{R}_q^p := R_q^p / (R_q^{p-1} + R_{q-1}^p) \quad \text{and} \quad \overline{R} = \overline{R}(\Delta) := \bigoplus_{p,q} \overline{R}_q^p.$$

The vector space $\overline{R}_q^p$ can be considered, roughly speaking, as the space of the homogeneous rational functions expressible as sums of fractions with numerators of degree p and denominators of degree q where p and q are the smallest possible. Since each $\overline{R}_q^p$ is finite-dimensional, one can define the **Poincaré series**

$$\text{Poin}(\overline{R}(\Delta), s, t) := \sum_{p,q} \dim(\overline{R}_q^p) s^p t^q$$

of the bigraded vector space $\overline{R}(\Delta)$. Let

$$\mathcal{A}(\Delta) = \{\ker(\alpha) \mid \alpha \in \Delta\}.$$

Then $\mathcal{A}(\Delta)$ is a (central) arrangement of hyperplanes [3] in V . Let $\text{Poin}(\mathcal{A}(\Delta), \cdot)$ be the Poincaré polynomial [3, Definition 2.48] of the arrangement $\mathcal{A}(\Delta)$, which is combinatorially defined. (See Definition 4.1.) Our main theorem is the following

Theorem 1.2. *We have*

$$\text{Poin}(\overline{R}(\Delta), s, t) = \frac{1}{(1-s)^\ell} \text{Poin}\left(\mathcal{A}(\Delta), \frac{t(1-s)}{(1-t)}\right).$$

By combining Theorem 1.2 and a later result (Proposition 2.2), one has

Corollary 1.3.

$$\sum_{p,q} \dim(R_q^p) s^p t^q = (1-s)^{-\ell-1} (1-t)^{-1} \text{Poin}\left(\mathcal{A}(\Delta), \frac{t(1-s)}{(1-t)}\right).$$

By Theorem 1.2 and the factorization theorem (Theorem 4.2) in [7], we may easily show the following

Corollary 1.4. *If $\mathcal{A}(\Delta)$ is a free arrangement with exponents $(d_1, \dots, d_\ell)$ [7] [3, Definitions 4.15, 4.25], then*

$$\text{Poin}(\overline{R}(\Delta), t) = (1-s)^{-\ell} (1-t)^{-\ell} \prod_{i=1}^{\ell} \{1 + (d_i - 1 - s d_i) t\}.$$

In particular, one has

Corollary 1.5. *When $\mathcal{A}(\Delta)$ is the set of reflecting hyperplanes of any (real or complex) reflection group with exponents $(d_1, \dots, d_\ell)$, [3, Definitions 6.22], then*

$$\text{Poin}(\overline{R}(\Delta), t) = (1-s)^{-\ell}(1-t)^{-\ell} \prod_{i=1}^{\ell} \{1 + (d_i - 1 - sd_i)t\}.$$

Example 1.6. Let $x_1, \dots, x_\ell$ be a basis for V^* . Let $\Delta = \{x_i - x_j \mid 1 \leq i < j \leq \ell\}$. Then Δ gives the root system of type $A_{\ell-1}$. The corresponding reflection arrangement $\mathcal{A}(\Delta)$ is a braid arrangement with exponents $(0, 1, \dots, \ell-1)$ [3, Example 4.32]. So, by Corollary 1.5, we have

$$\begin{aligned} \text{Poin}(\overline{R}(\Delta), s, t) &= (1-s)^{-\ell}(1-t)^{-\ell+1} \prod_{i=1}^{\ell-1} \{1 + (i-1)t - ist\} \\ &= (1-s)^{-\ell}(1-t)^{-\ell+1}(1-st)(1+t-2st) \dots (1+(\ell-2)t - (\ell-1)st). \end{aligned}$$

For instance, when $\ell = 3$, we have

$$R(\Delta) = \mathbf{K} \left[x_1, x_2, x_3, \frac{1}{x_1 - x_2}, \frac{1}{x_2 - x_3}, \frac{1}{x_1 - x_3} \right]$$

and

$$\begin{aligned} &\text{Poin}(\overline{R}(\Delta), s, t) \\ &= \frac{(1-st)(1+t-2st)}{(1-s)^3(1-t)^2} \\ &= \sum_{p \geq 0} \binom{p+2}{2} s^p + \sum_{p \geq 0} \sum_{q > 0} (3p+2q+1) s^p t^q \\ &= 1 + (3s+3t) + (6s^2+6st+5t^2) + (10s^3+9s^2t+8st^2+7t^3) \\ &\quad + (15s^4+12s^3t+11s^2t^2+10st^3+9t^4) + \dots \end{aligned}$$

Since the coefficient of st^2 is equal to 8, the space $\overline{R}_2^1$ is eight-dimensional. Actually we can easily verify by direct computation that the classes of

$$\begin{aligned} &\frac{x_1}{(x_1-x_2)(x_1-x_3)}, \frac{x_1}{(x_1-x_2)(x_2-x_3)}, \frac{x_1}{(x_1-x_2)^2}, \frac{x_3}{(x_1-x_2)^2}, \\ &\frac{x_1}{(x_2-x_3)^2}, \frac{x_2}{(x_2-x_3)^2}, \frac{x_1}{(x_1-x_3)^2}, \frac{x_2}{(x_1-x_3)^2} \end{aligned}$$

form a basis for $\overline{R}_2^1$. This implies that an arbitrary element φ of R_2^1 can be uniquely expressed as a $\mathbf{K}$ -linear combination of these eight rational functions modulo $R_2^0 + R_1^1$. For example, when

$$\varphi = \frac{x_1 + 2x_2 - x_3 + 3}{(x_1 - x_3)(x_2 - x_3)}$$

φ is decomposed into partial fractions as

$$\begin{aligned} \varphi = & -2 \frac{x_1}{(x_1 - x_2)(x_1 - x_3)} + 2 \frac{x_1}{(x_1 - x_2)(x_2 - x_3)} + \frac{3}{(x_1 - x_3)(x_2 - x_3)} \\ & - \frac{1}{x_2 - x_3} + \frac{2}{x_1 - x_3}, \end{aligned}$$

which implies

$$\varphi \equiv -2 \frac{x_1}{(x_1 - x_2)(x_1 - x_3)} + 2 \frac{x_1}{(x_1 - x_2)(x_2 - x_3)} \pmod{R_2^0 + R_1^1}.$$

Remark 1.7. The specializations of $\text{Poin}(\overline{R}(\Delta), s, t)$ at $s = 0$ and $t = 0$ have been known:

(1) Suppose $t = 0$. Since $\overline{R}_0^p = R_0^p/R_0^{p-1}$ is isomorphic to the vector space of homogeneous polynomials of degree p in ℓ variables, we have $\bigoplus_p \overline{R}_0^p \simeq S$ and thus

$$\text{Poin}(\overline{R}(\Delta), s, 0) = \frac{1}{(1-s)^\ell}.$$

(2) Suppose $s = 0$. Note that $\overline{R}_q^0 = R_q^0/R_{q-1}^0$ is isomorphic to a $\mathbf{K}$ -vector subspace

$$C(\Delta)_q := \left\langle \frac{1}{g} \in R \mid g \text{ is a product of } q \text{ linear forms in } \Delta \right\rangle_{\mathbf{K}}$$

of $R(\Delta)$. The $\mathbf{K}$ -subalgebra

$$C(\Delta) := \bigoplus_q C(\Delta)_q$$

of $R(\Delta)$ was studied and the formula

$$\text{Poin}(C(\Delta), t) = \sum_{q \geq 0} (\dim C(\Delta)_q) t^q = \text{Poin}(\mathcal{A}(\Delta), (1-t)^{-1}t)$$

was obtained in [7, Theorem 1.4]. (The proof of this formula in [7] heavily depends on [1] and [4].) So we have

$$\text{Poin}(\overline{R}(\Delta), 0, t) = \text{Poin}(\mathcal{A}(\Delta), (1-t)^{-1}t).$$

2 The bigraded vector space $\overline{R}(\Delta)$

Let $p, q \in \mathbb{Z}$. Let $\pi_q^p : R_q^p \rightarrow \overline{R}_q^p$ be the canonical projection. Fix a linear section

$$\iota_q^p : \overline{R}_q^p \longrightarrow R_q^p$$

for each canonical projection π_q^p such that

$$\text{Image}(\iota_q^p) \subseteq \left\langle \frac{f}{g} \in R \mid f \text{ and } g \text{ are homogeneous with } \deg f = p \text{ and } \deg g = q \right\rangle_{\mathbf{K}}.$$

Proposition 2.1. *There exists a linear isomorphism*

$$\kappa := \sum_{p,q} \iota_q^p : \overline{R} = \bigoplus_{p,q} \overline{R}_q^p \xrightarrow{\sim} \sum_{p,q} R_q^p = R.$$

Proof. At first, we prove the injectivity of κ . For every $\phi = \sum_{p,q} \phi_q^p \in \ker(\kappa)$, where $\phi_q^p \in \overline{R}_q^p$, one has

$$0 = \kappa(\phi) = \sum_{p,q} \iota_q^p(\phi_q^p) = \sum_{a \in \mathbb{Z}} \sum_{p-q=a} \iota_q^p(\phi_q^p).$$

This implies that $\sum_{p-q=a} \iota_q^p(\phi_q^p) = 0$ for each $a \in \mathbb{Z}$ because $\deg \iota_q^p(\phi_q^p) = p - q$. Suppose $\phi \neq 0$. Then there exists an integer $b \in \mathbb{Z}$ such that $\sum_{p-q=b} \phi_q^p \neq 0$. Let $p_b := \max\{p \mid \phi_q^p \neq 0, p - q = b\}$ and $q_b = p_b - b$. Then $\psi_{q_b}^{p_b} := \iota_{q_b}^{p_b}(\phi_{q_b}^{p_b}) \neq 0$, because $\iota_{q_b}^{p_b}$ is injective. Therefore one has

$$0 = \sum_{p-q=b} \iota_q^p(\phi_q^p) = \psi_{q_b}^{p_b} + \psi_{q_b-1}^{p_b-1} + \dots$$

and

$$\psi_{q_b}^{p_b} = -\psi_{q_b-1}^{p_b-1} - \psi_{q_b-2}^{p_b-2} - \dots \in R_{q_b-1}^{p_b-1} \subseteq R_{q_b-1}^{p_b} + R_{q_b}^{p_b-1}.$$

So we obtain

$$\phi_{q_b}^{p_b} = \pi_{q_b}^{p_b}(\iota_{q_b}^{p_b}(\phi_{q_b}^{p_b})) = \pi_{q_b}^{p_b}(\psi_{q_b}^{p_b}) = 0,$$

which contradicts the definition of p_b . Therefore we have $\ker(\kappa) = \{0\}$.

Next we prove the surjectivity of κ . Let $p \geq 0$ and $q \geq 0$. Define

$$\kappa_q^p := \sum_{a \leq p, b \leq q} \iota_b^a : \bigoplus_{a \leq p, b \leq q} \overline{R}_b^a \longrightarrow R_q^p.$$

It is sufficient to show that κ_q^p is surjective for $p, q \in \mathbb{Z}$. If either p or q is negative, then $R_q^p = 0$ and there is nothing to prove. We will prove $R_q^p = \text{Image}(\kappa_q^p)$ for $p + q \geq 0$ by an induction on $p + q$. When $p = q = 0$, $\pi_0^0 : \mathbf{K} = R_0^0 \rightarrow \overline{R}_0^0 = \mathbf{K}$ is an isomorphism. So $R_0^0 = \text{Image}(\iota_0^0) = \text{Image}(\kappa_0^0)$. Suppose $p + q > 0$. For $\psi \in R_q^p$, let $\eta := \psi - \iota_q^p \circ \pi_q^p(\psi)$. Then

$$\pi_q^p(\eta) = \pi_q^p(\psi) - \pi_q^p \circ \iota_q^p \circ \pi_q^p(\psi) = 0.$$

So $\eta \in \ker \pi_q^p = R_q^{p-1} + R_{q-1}^p \subseteq \text{Image}(\kappa_q^p)$ by the induction assumption. We thus have

$$\psi = \eta + \iota_q^p \circ \pi_q^p(\psi) \in \text{Image}(\kappa_q^p),$$

which implies that κ_q^p is surjective. $\square$

In the course of the proof of Proposition 2.1, we have already proved the following

Proposition 2.2. *Let $p \geq 0$ and $q \geq 0$. Then there exists a linear isomorphism*

$$\kappa_q^p = \sum_{a \leq p, b \leq q} \iota_b^a : \bigoplus_{a \leq p, b \leq q} \overline{R}_b^a \xrightarrow{\sim} R_q^p.$$

3 A decomposition of $\overline{R_q^p}$

Let $\mathbf{E}_p(\Delta)$ be the set of all p -tuples composed of elements of Δ . Let $\mathbf{E}(\Delta) := \bigcup_{p \geq 0} \mathbf{E}_p(\Delta)$. The union is disjoint. Write $\prod \mathcal{E} := \alpha_1 \dots \alpha_p \in S$ when $\mathcal{E} = (\alpha_1, \dots, \alpha_p) \in \mathbf{E}_p(\Delta)$. For $\varepsilon \in \mathbf{E}(\Delta)$, let $V(\varepsilon)$ denote the set of common zeros of ε :

$$V(\varepsilon) = \bigcap_{i=1}^p \ker(\alpha_i)$$

when $\varepsilon = (\alpha_1, \dots, \alpha_p)$. Define

$$L = L(\Delta) = \{V(\varepsilon) \mid \varepsilon \in \mathbf{E}(\Delta)\}.$$

Agree that $V(\varepsilon) = V$ if ε is the empty tuple. Introduce a partial order $\leq$ into L by reverse inclusion: $X \leq Y \Leftrightarrow X \supseteq Y$. Then L is equal to the intersection lattice of the arrangement $\mathcal{A}(\Delta)$ [3, Definition 2.1]. For $X \in L$, define

$$\mathbf{E}_X(\Delta) := \{\varepsilon \in \mathbf{E}(\Delta) \mid V(\varepsilon) = X\}.$$

Then

$$\mathbf{E}(\Delta) = \bigcup_{X \in L} \mathbf{E}_X(\Delta) \quad (\text{disjoint}).$$

Define

$$C_X(\Delta) := \sum_{\varepsilon \in \mathbf{E}_X(\Delta)} \mathbf{K}(\prod \varepsilon)^{-1}.$$

The following proposition is in [7, Proposition 2.1]:

Proposition 3.1.

$$C(\Delta) = \bigoplus_{X \in L} C_X(\Delta).$$

Let $X \in L$ and $q \in \mathbb{Z}_{\geq 0}$. Define

$$C_{q,X} := C_{q,X}(\Delta) := C(\Delta)_q \cap C_X(\Delta) = \sum_{\varepsilon \in \mathbf{E}_q(\Delta) \cap \mathbf{E}_X(\Delta)} \mathbf{K}(\prod \varepsilon)^{-1}.$$

Then

$$C_X(\Delta) = \bigoplus_{q \geq 0} C_{q,X}.$$

Definition 3.2. ([3, 2.42]) Let the **Möbius function**

$$\mu : L(\Delta) \rightarrow \mathbb{Z}$$

be characterized by $\mu(V) = 1$ and for $X < V$ by $\sum_{Y \leq X} \mu(Y) = 0$.

The following result is in [7, Theorem 1.4]:

Proposition 3.3.

$$\text{Poin}(C_X(\Delta), t) := \sum_q (\dim C_{q,X}) t^q = (-1)^{\text{codim} X} \mu(X) \left(\frac{t}{1-t} \right)^{\text{codim} X}.$$

Define $I(X) := \{f \in S \mid f|_X \equiv 0\}$, i.e., $I(X)$ is the prime ideal of S generated by the polynomial functions vanishing on X . Let $S_X := S/I(X)$. Let S^p denote the degree p homogeneous part of S . Then $S = \bigoplus_{p \geq 0} S^p$. Define

$$S_X^p := S^p / S^p \cap I(X).$$

Then

$$S_X := \bigoplus_{p \geq 0} S_X^p.$$

Lemma 3.4. *For $p \geq 0$, $q \geq 0$, and $X \in L$, suppose that $h_1, h_2, \dots, h_m \in C_{q,X}$ are linearly independent over $\mathbf{K}$ and $f_1, f_2, \dots, f_m \in S^p$. Then the following three conditions are equivalent:*

- (1) *Each f_i belongs to the ideal $I(X)$,*
- (2) $\sum_{i=1}^m f_i h_i \in R_{q-1}^{p-1}$,
- (3) $\sum_{i=1}^m f_i h_i \in R_{q-1}^{p-1} + \sum_{\substack{Y \in L \\ Y \neq X}} S^p C_{q,Y}$.

Proof. (1) $\Rightarrow$ (2) : If $f_i \in S^p \cap I(X)$, then one can easily see that $f_i h_i \in R_{q-1}^{p-1}$ for each i .

(2) $\Rightarrow$ (3) : Obvious.

(3) $\Rightarrow$ (1) : Let

$$\prod \Delta = \prod_{\alpha \in \Delta} \alpha, \quad \Delta_X = \Delta \cap I(X), \quad \text{and} \quad \prod \Delta_X = \prod_{\alpha \in \Delta_X} \alpha.$$

Suppose that $\dim X = k$. Choose a basis $x_1, x_2, \dots, x_\ell$ for V^* such that $X = V(x_{k+1}, \dots, x_\ell)$. Let

$$h'_i := \left(\prod \Delta_X \right)^q h_i \in \mathbf{K}[x_{k+1}, \dots, x_\ell],$$

which is a homogeneous polynomial of degree $q|\Delta_X| - q$. Multiply $(\prod \Delta_X)^q$ to $\sum_{i=1}^m f_i h_i \in R_{q-1}^{p-1} + \sum_{\substack{Y \in L \\ Y \neq X}} S^p C_{q,Y}$ to get

$$\left(\frac{\prod \Delta}{\prod \Delta_X} \right)^q \sum_i f_i h'_i = \left(\prod \Delta \right)^q \sum_i f_i h_i \in I(X)^{q|\Delta_X| - q + 1}.$$

Since $I(X)^{q|\Delta_X| - q + 1}$ is primary and $\prod \Delta / \prod \Delta_X \notin I(X)$, one has

$$\sum_i f_i h'_i \in I(X)^{q|\Delta_X| - q + 1} = (x_{k+1}, \dots, x_\ell)^{q|\Delta_X| - q + 1}.$$

Define $f'_i := f_i(x_1, \dots, x_k, 0, \dots, 0) \in \mathbf{K}[x_1, \dots, x_k]$. Since $f_i - f'_i \in I(X)$, one has

$$\sum_i f'_i h'_i \in I(X)^{q|\Delta_X| - q + 1} = (x_{k+1}, \dots, x_\ell)^{q|\Delta_X| - q + 1}.$$

Recalling that each f'_i lies in $\mathbf{K}[x_1, \dots, x_k]$ and $\deg h'_i = q|\Delta_X| - q$ with $h'_i \in (x_{k+1}, \dots, x_\ell)^{q|\Delta_X| - q}$, we may conclude

$$\sum_i f'_i h'_i = 0.$$

We may deduce that $f'_1 = f'_2 = \dots = f'_m = 0$ because $h'_1, \dots, h'_m \in \mathbf{K}[x_{k+1}, \dots, x_\ell]$ are linearly independent over $\mathbf{K}$ and thus over $\mathbf{K}[x_1, \dots, x_k]$ also. Therefore $f_i \in I(X)$ for each i . $\square$

Proposition 3.5. *For $p \geq 0$, $q \geq 0$, and $X \in L$, let*

$$\tau_{q,X}^p : S_X^p \otimes C_{q,X} \longrightarrow \overline{R_q^p}$$

be the linear map characterized by

$$\tau_{q,X}^p \left([f] \otimes (\prod \varepsilon)^{-1} \right) = \left[\frac{f}{(\prod \varepsilon)} \right],$$

where $f \in S^p$ and $\varepsilon \in \mathbf{E}_q(\Delta) \cap \mathbf{E}_X(\Delta)$. Then $\tau_{q,X}^p$ is injective for each $X \in L$.

Proof. Let $\varepsilon \in \mathbf{E}_q(\Delta) \cap \mathbf{E}_X(\Delta)$. Note that ε generates the ideal $I(X)$. If f belongs to $S^p \cap I(X)$, then

$$\frac{f}{(\prod \varepsilon)} \in R_{q-1}^{p-1} \subset R_q^{p-1} + R_{q-1}^p$$

by Lemma 3.4. This implies that the map $\tau_{q,X}^p$ is well-defined.

Choose a $\mathbf{K}$ -basis $h_1, h_2, \dots, h_m$ for $C_{q,X}$. Then an arbitrary element $\varphi \in S_X^p \otimes C_{q,X}$ can be expressed as

$$\varphi = \sum_{i=1}^m [f_i] \otimes h_i$$

for some $f_i \in S^p$ ($i = 1, \dots, m$). Suppose $\varphi \in \ker(\tau_{q,X}^p)$, i.e., $\sum_{i=1}^m f_i h_i \in R_q^{p-1} + R_{q-1}^p$. This implies $\sum_{i=1}^m f_i h_i \in R_{q-1}^{p-1}$. By Lemma 3.4, one has $f_i \in I(X)$ for each i . Therefore $\varphi = 0$ and thus $\tau_{q,X}^p$ is injective. $\square$

Proposition 3.6. *Define $\overline{R_{q,X}^p} := \text{Image}(\tau_{q,X}^p)$. Then we have*

$$\overline{R_q^p} = \bigoplus_{X \in L} \overline{R_{q,X}^p}.$$

Proof. An arbitrary element $\varphi \in \overline{R_q^p}$ can be expressed as

$$\varphi = \left[\sum_i \frac{f_i}{g_i} \right]$$

with $\deg(f_i) = p$ and $\deg(g_i) = q$. Thus we have

$$\overline{R_q^p} \subseteq \sum_{X \in L} \overline{R_{q,X}^p}.$$

Suppose that $\sum_{X \in L} \phi_X = 0$ in $\overline{R_q^p}$ where $\phi_X \in \overline{R_{q,X}^p}$. We will prove $\phi_X = 0$ for each X . Fix $X \in L$. Choose a $\mathbf{K}$ -basis $h_1, h_2, \dots, h_m$ for $C_{q,X}$. Write

$$\phi_X = \tau_{q,X}^p \left(\sum_i [f_i] \otimes h_i \right) = \left[\sum_i f_i h_i \right]$$

for some $f_i \in S^p$ for each i . Since

$$\left[\sum_i f_i h_i \right] = \phi_X = - \sum_{\substack{Y \in L \\ Y \neq X}} \phi_Y,$$

we have

$$\sum_i f_i h_i \in R_{q-1}^{p-1} + \sum_{\substack{Y \in L \\ Y \neq X}} S^p C_{q,Y}.$$

By Lemma 3.4, one has $f_i \in I(X)$ for each i . Thus

$$\phi_X = \tau_{q,X}^p \left(\sum_i [f_i] \otimes h_i \right) = 0. \quad \square$$

4 Proofs

In this section we prove Theorem 1.2, Corollaries 1.4 and 1.5. First recall the Möbius function $\mu : L \rightarrow \mathbb{Z}$ in Definition 3.2.

Definition 4.1. *The Poincaré polynomial of $\mathcal{A}(\Delta)$ is defined by*

$$\text{Poin}(\mathcal{A}(\Delta), t) = \sum_{X \in L} \mu(X) (-t)^{\text{codim} X}.$$

The Poincaré polynomial is combinatorially defined and is known to be equal to the Poincaré polynomial of the topological space $M(\mathcal{A}(\Delta)) := V \setminus \bigcup_{H \in \mathcal{A}(\Delta)} H$ when $\mathbf{K} = \mathbb{C}$ [2] [3, Theorem 5.93].

Proof of Theorem 1.2. By Propositions 3.6 and 3.5, we have

$$\overline{R_q^p} = \bigoplus_{X \in L} \overline{R_{q,X}^p} \quad \text{and} \quad \overline{R_{q,X}^p} \cong S_X^p \otimes C_{q,X}.$$

Combining these with Proposition 3.3, we have

$$\begin{aligned} \text{Poin}(\overline{R}(\Delta), s, t) &= \sum_{p,q} \dim(\overline{R_q^p}) s^p t^q = \sum_{p,q} \left(\sum_{X \in L} \dim(\overline{R_{q,X}^p}) \right) s^p t^q \\ &= \sum_{X \in L} \left(\sum_{p \geq 0} \dim(S_X^p) s^p \right) \left(\sum_{q \geq 0} \dim(C_{q,X}) t^q \right) \\ &= \sum_{X \in L} \left(\frac{1}{1-s} \right)^{\dim X} (-1)^{\text{codim} X} \mu(X) \left(\frac{t}{1-t} \right)^{\text{codim} X} \\ &= \sum_{X \in L} (-1)^{\text{codim} X} \mu(X) \left(\frac{t(1-s)}{1-t} \right)^{\text{codim} X} \left(\frac{1}{1-s} \right)^\ell \\ &= \frac{1}{(1-s)^\ell} \text{Poin} \left(\mathcal{A}(\Delta), \frac{t(1-s)}{1-t} \right). \quad \square \end{aligned}$$

Proof of Corollary 1.3. By Propostion 2.2, we obtain

$$\dim(R_q^p) = \sum_{a \leq p, b \leq q} \dim(\overline{R_b^a}).$$

Thus

$$\begin{aligned} \sum_{p,q} \dim(R_q^p) s^p t^q &= \left(\sum_{c \geq 0} s^c \right) \left(\sum_{d \geq 0} t^d \right) \sum_{a,b} \dim(\overline{R_b^a}) s^a t^b \\ &= \frac{1}{(1-s)(1-t)} \text{Poin}(\overline{R}(\Delta), s, t) = \frac{1}{(1-s)^{\ell+1}(1-t)} \text{Poin} \left(\mathcal{A}(\Delta), \frac{t(1-s)}{1-t} \right). \quad \square \end{aligned}$$

Let Der be the S -module of derivations :

$$\text{Der} = \{ \theta \mid \theta : S \rightarrow S \text{ is a } \mathbf{K}\text{-linear derivations} \}.$$

Then Der is naturally isomorphic to $S \otimes_{\mathbf{K}} V$. Define

$$D(\Delta) = \{ \theta \in \text{Der} \mid \theta(\alpha) \in \alpha S \text{ for any } \alpha \in \Delta \},$$

which is naturally an S -submodule of Der. We say that the arrangement $\mathcal{A}(\Delta)$ is **free** if $D(\Delta)$ is a free S -module [3, Definition 4.15]. An element $\theta \in D(\Delta)$ is said to be **homogeneous of degree p** if

$$\theta(x) \in S^p \text{ for all } x \in V^*.$$

When $\mathcal{A}(\Delta)$ is a free arrangement, let $\theta_1, \dots, \theta_\ell$ be a homogeneous basis for $D(\Delta)$. The ℓ nonnegative integers $\deg \theta_1, \dots, \deg \theta_\ell$ are called the **exponents** of $\mathcal{A}(\Delta)$. Then one has

Proposition 4.2. (Factorization Theorem [7], [3, Theorem 4.137]) *If $\mathcal{A}(\Delta)$ is a free arrangement with exponents $d_1, \dots, d_\ell$, then*

$$\text{Poin}(\mathcal{A}(\Delta), t) = \prod_{i=1}^{\ell} (1 + d_i t).$$

By Theorem 1.2 and Proposition 4.2, we immediately have Corollary 1.4. Corollary 1.5 is a special case of Corollary 1.4 because the set $\mathcal{A}(\Delta)$ of reflecting hyperplanes is known to be a free arrangement [5] [6] [3, Theorem 6.60].

References

- [1] Brion, M., Vergne, M.: Arrangement of hyperplanes I. Rational functions and Jeffrey-Kirwan residue. Ann. scient. Éc. Norm. Sup., **32** (1999) 715–741
- [2] Orlik, P., Solomon, L.: Combinatorics and topology of complements of hyperplanes. Inventiones math. **56** (1980), 167–189
- [3] Orlik, P., Terao, H.: *Arrangements of Hyperplanes*. Grundlehren der Math. Wiss. **300**, Springer Verlag, 1992
- [4] Orlik, P., Terao, H.: Commutative algebras for arrangements, Nagoya J. Math., **134** (1994) 65 – 73
- [5] Saito, K.: On the uniformization of complements of discriminant loci. In: Conference Notes. Amer. Math. Soc. Summer Institute, Williamstown, 1975
- [6] Terao, H.: Free arrangements of hyperplanes and unitary reflection groups. Proc. Japan Acad. Ser. A **56** (1980) 389–392
- [7] Terao, H.: Generalized exponents of a free arrangement of hyperplanes and Shepherd-Todd-Brieskorn formula. Inventiones math. **63** (1981), 159–179
- [8] Terao, H.: Algebras generated by reciprocals of linear forms, to appear in J.Algebra